

On deformations of $\mathbf{Q}$ -factorial symplectic varieties

Yoshinori Namikawa

1 Introduction

This is a supplement to [Na]. In [Na] we have proposed a new category of complex symplectic varieties which admit certain singularities. It includes symplectic V-manifolds (cf. [Fu 1]) and O’Grady’s singular moduli spaces of semi-stable torsion free sheaves on a K3 surface [O] (cf. [Na], Introduction). A normal compact Kähler space Z is a symplectic variety if its regular locus U admits a non-degenerate holomorphic closed 2-form ω with the following property: for any resolution $\nu : \tilde{Z} \rightarrow Z$ of Z with $\nu^{-1}(U) \cong U$, the 2-form ω on $\nu^{-1}(U)$ extends to a holomorphic 2-form on $\tilde{Z}$. If Z can be embedded in a projective space, Z is called a projective symplectic variety.

Our purpose is to give a positive answer to the following problem posed in [Na]:

Problem. *Let Z be a $\mathbf{Q}$ -factorial projective symplectic variety with terminal singularities. Assume that Z is smoothable by a suitable flat deformation. Is Z then non-singular from the first ?*

We actually prove more:

Main Theorem. *Let Z be a $\mathbf{Q}$ -factorial projective symplectic variety with terminal singularities. Then any flat deformation of Z is locally trivial; in other words, it preserves all singularities on Z .*

This theorem would explain why “local Torelli theorem” [Na, Theorem 8] holds for such symplectic varieties.

Example ([O], cf. also [K-L-S]): Let S be a K3 surface with Picard number 1. Fix an even integer c with $c \geq 6$. Let M_c be the moduli space of semi-stable sheaves on S of rank 2 with $c_1 = 0$ and $c_2 = c$. O’Grady proved in

[O] that M_c is a projective symplectic variety with terminal singularities. In the preprint version of [O]: math.AG/9708009, he constructed a projective resolution $\hat{\pi} : \hat{M}_c \rightarrow M_c$. Let $N_1(\hat{M}_c/M_c)$ be the $\mathbf{R}$ -vector space of $\hat{\pi}$ -numerical classes of 1-cycles l on $\hat{M}_c$ such that $\hat{\pi}(l) = \text{point}$. Then

$$N_1(\hat{M}_c/M_c) = \mathbf{R}[\hat{e}_c] \oplus \mathbf{R}[\hat{\sigma}_c] \oplus \mathbf{R}[\hat{\gamma}_c]$$

for effective 1-cycles $\hat{e}_c$, $\hat{\sigma}_c$ and $\hat{\gamma}_c$ (ibid. Section 3). On the other hand, $\text{Exc}(\hat{\pi})$ consists of three prime divisors $\hat{\Omega}_c$, $\hat{\Sigma}_c$ and $\hat{\Delta}_c$ (ibid. Section 3). This implies that M_c is $\mathbf{Q}$ -factorial. In fact, let D be a prime divisor of M_c and let D' be its proper transform on $\hat{M}_c$. By the observation above, we can find $a_i \in \mathbf{Q}$ ($i = 1, 2, 3$) such that $D'' := D' + a_1\hat{\Omega}_c + a_2\hat{\Sigma}_c + a_3\hat{\Delta}_c$ is $\hat{\pi}$ -numerically trivial. Since M_c has only rational singularities, there is a positive integer m such that mD'' is linearly equivalent to the pull-back of a Cartier divisor E of M_c . Then mD is linearly equivalent to E . More generally, for a K3 surface or an Abelian surface, almost all moduli spaces of semi-stable sheaves with non-primitive Mukai vectors are $\mathbf{Q}$ -factorial projective symplectic varieties [K-L-S]. Applying Main Theorem to these moduli spaces, we conclude that any deformation of them are locally rigid; in particular, they have no smoothing via deformations.

In the final section, we shall apply Main Theorem to get the following remarkable results.

Corollary 1. *Let $h : Y \rightarrow Z$ and $h' : Y' \rightarrow Z$ be two $\mathbf{Q}$ -factorial terminalizations of a projective symplectic variety Z ; in other words, h (resp. h') is a crepant projective birational morphism such that Y (resp. Y') has only $\mathbf{Q}$ -factorial terminal singularities. Assume that Y is non-singular. Then Y' is also non-singular.*

Remark 1. A $\mathbf{Q}$ -factorial terminalization Y of a projective symplectic variety Z is again symplectic. In fact, there is a 2-form $h^*\omega$ on $h^{-1}(U)$, where $U := Z_{\text{reg}}$. By the definition of a symplectic variety, this 2-form extends to a holomorphic 2-form on $V := Y_{\text{reg}}$. Moreover, since $K_Y = h^*K_Z$, the extended 2-form is non-degenerate over V . If we take a resolution $\tilde{Y}$ of Y , then this 2-form further extends to a holomorphic 2-form on $\tilde{Y}$.

Example (flops): When the codimensions of $\text{Exc}(h)$ and $\text{Exc}(h')$ are both larger than one, Corollary 1 says that *the smoothness is preserved under a flop*. A similar result has been proved by Kaledin under certain conditions when Z is an affine symplectic singularity (cf. “Symplectic resolutions:

deformations and birational maps”, math.AG/0012008).

Corollary 2. *Let Z be a projective symplectic variety. Assume that Z has a $\mathbf{Q}$ -factorial terminalization $h : Y \rightarrow Z$. Then the following are equivalent.*

- (1) Z is smoothable by a flat deformation;
- (2) Y is non-singular.

The following conjecture would be true if any projective symplectic variety had a $\mathbf{Q}$ -factorial terminalization (cf. Remark 2, (1))

Conjecture. *A projective symplectic variety is smoothable by a deformation if and only if it admits a crepant (symplectic) resolution.*

Remark 2. (1) The existence of a $\mathbf{Q}$ -factorial terminalization follows from the “minimal model conjecture”. In fact, let $\tilde{Z} \rightarrow Z$ be a projective resolution. If the minimal model conjecture holds, one can get a $\mathbf{Q}$ -factorial terminalization $Y \rightarrow Z$ after running the minimal model program for $\tilde{Z}$ over the fixed base Z . The $\mathbf{Q}$ -factorial terminalizations are not unique; however, they have the same kind of singularities by virtue of the proof of Corollary 1. The minimal model conjecture is proved by Shokurov [Sh] in dimension 4.

(2) It would be interesting to know whether our results also hold in the case where Z is a *convex* symplectic variety. Here we say that Z is *convex* if there is a birational projective morphism from Z to an affine symplectic singularity X . When X has non-isolated singularity, the usual deformation theory of Z (and X) has an infinite dimensional nature and does not work well. Instead, we should consider a “Poisson deformation” (cf. [G-K]).

2 Proof of Main Theorem

(i) **$\mathbf{Q}$ -factoriality:** A normal variety Z is called $\mathbf{Q}$ -factorial if, for any Weil divisor D of Z , mD is a Cartier divisor of Z for some positive integer m . Assume that Z is a projective variety with rational singularities. Let $\nu : \tilde{Z} \rightarrow Z$ be a resolution such that the exceptional locus consists of finite number of divisors E_i ($1 \leq i \leq m$). Denote by $[E_i] \in H^2(\tilde{Z}, \mathbf{Z})$ the corresponding classes. By [Ko-Mo, (12.1.6)], Z is $\mathbf{Q}$ -factorial if and only if

$$\mathrm{im}[H^2(\tilde{Z}, \mathbf{Q}) \rightarrow H^0(Z, R^2\nu_*\mathbf{Q})] = \mathrm{im}[\oplus \mathbf{Q}[E_i] \rightarrow H^0(Z, R^2\nu_*\mathbf{Q})].$$

We give here a proof of the “only if” part. Fix an ample class $\kappa \in H^2(\tilde{Z}, \mathbf{Q})$. Let $H_0^2(\tilde{Z}, \mathbf{Q})$ be the primitive part of $H^2(\tilde{Z}, \mathbf{Q})$. By definition, we have

$$H^2(\tilde{Z}, \mathbf{Q}) = H_0^2(\tilde{Z}, \mathbf{Q}) \oplus \mathbf{Q}\kappa.$$

Since Z has only rational singularities, the natural map $H^2(Z, \mathbf{Q}) \rightarrow H^2(\tilde{Z}, \mathbf{Q})$ is an injection of mixed Hodge structures ([De]); hence, by the strict compatibility of weight filtrations, $H^2(Z, \mathbf{Q})$ admits a pure $\mathbf{Q}$ -Hodge structure of weight 2. Then $H_0^2(Z, \mathbf{Q}) := H^2(Z, \mathbf{Q}) \cap H_0^2(\tilde{Z}, \mathbf{Q})$ also admits a pure $\mathbf{Q}$ -Hodge structure of weight 2. Note that $H_0^2(Z, \mathbf{C})$ contains $H^{2,0}(\tilde{Z})$ and $H^{0,2}(\tilde{Z})$. With respect to the Hodge-Riemann bilinear form on $H_0^2(\tilde{Z}, \mathbf{Q})$, we put $V := (H_0^2(Z, \mathbf{Q}))^\perp$. One can check that

$$V \cap H_0^2(Z, \mathbf{Q}) = 0,$$

and hence

$$H^2(\tilde{Z}, \mathbf{Q}) = V \oplus \mathbf{Q}\kappa \oplus H_0^2(Z, \mathbf{Q}).$$

Note that $H_0^2(Z, \mathbf{Q})$ is mapped to zero by the map $H^2(\tilde{Z}, \mathbf{Q}) \rightarrow H^0(Z, R^2\nu_*\mathbf{Q})$. Since every element of V is of type (1,1), we see that

$$\mathrm{im}[H^2(\tilde{Z}, \mathbf{Q}) \rightarrow H^0(Z, R^2\nu_*\mathbf{Q})] = \mathrm{im}[\mathrm{Pic}(\tilde{Z}) \otimes \mathbf{Q} \rightarrow H^0(Z, R^2\nu_*\mathbf{Q})].$$

Now take a $\mathbf{Q}$ -Cartier divisor D of $\tilde{Z}$. The push-forward ν_*D is a $\mathbf{Q}$ -Cartier divisor of Z since Z is $\mathbf{Q}$ -factorial. Now the pull-back $\nu^*(\nu_*D)$ of ν_*D is well-defined, and one has

$$D = \nu^*(\nu_*D) + \sum a_i E_i$$

for some $a_i \in \mathbf{Q}$. This implies the “only if” part.

(ii) **Terminal singularity:** We recall the definition of a terminal singularity. Assume, for simplicity, that Z is a normal variety such that the canonical divisor K_Z is a Cartier divisor. We say that Z has only terminal (resp. canonical) singularities if, for a resolution $\nu : \tilde{Z} \rightarrow Z$ such that the exceptional locus consists of finite number of divisors E_i ($1 \leq i \leq m$),

$$K_{\tilde{Z}} = \nu^*K_Z + \sum a_i E_i$$

with $a_i > 0$ (resp. $a_i \geq 0$) for all i . Let $\Sigma \subset Z$ be the singular locus. If Z has only terminal singularities, then we have $\mathrm{codim}(\Sigma \subset Z) \geq 3$ (cf. [Re]). But

if, in addition, Z is a symplectic variety, then we have $\text{codim}(\Sigma \subset Z) \geq 4$ by [Na 2], [K]. Conversely, if Z is a symplectic variety with $\text{codim}(\Sigma \subset Z) \geq 4$, then Z has only terminal singularities [Na 2].

(iii) **Infinitesimal deformations:** Assume that Z is a reduced complex space. Recall that a deformation $\mathcal{Z}$ of Z over an analytic space S with a reference point $0 \in S$ is a flat map $\mathcal{Z} \rightarrow S$ with a fixed isomorphism $\mathcal{Z} \times_S \{0\} \cong Z$. When $0 \in S$ is a spectrum of an Artinian local $\mathbf{C}$ -algebra with a unique point, $\mathcal{Z}$ is called an infinitesimal deformation of Z . Moreover, an infinitesimal deformation $\mathcal{Z} \rightarrow S$ is called *locally trivial* if, for any Stein open set $V \subset Z$, $\mathcal{Z}|_V \cong V \times S$.

For $n \geq 0$, we put $A_n := \mathbf{C}[t]/(t^{n+1})$ and $S_n := \text{Spec} A_n$. Suppose that $Z_n \rightarrow S_n$ is an infinitesimal deformation of Z over S_n . Define $T_{Z_n/S_n} := \underline{\text{Hom}}(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n})$. Let $S_n[\epsilon] := \text{Spec}(A_n[\epsilon]/\epsilon^2)$. There is a natural closed immersion $S_n \rightarrow S_n[\epsilon]$. Moreover, the injection $A_n \rightarrow A_n[\epsilon]/\epsilon^2$ induces a map $S_n[\epsilon] \rightarrow S_n$. $\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n})$ plays an important role in the deformation theory of a *reduced* complex space:

(iii-a): *There is a one-to-one correspondence between $\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n})$ and the set of equivalence classes of infinitesimal deformations of Z_n over $S_n[\epsilon]$.*

Here two infinitesimal deformations $\mathcal{Z}_n$ and $\mathcal{Z}'_n$ of Z_n over $S_n[\epsilon]$ are equivalent if there is an $S_n[\epsilon]$ -isomorphism between them which induces, over S_n , the identity map of Z_n .

The correspondence is given in the following manner. Assume an infinitesimal deformation $\mathcal{Z}_n$ is given. By the map $S_n[\epsilon] \rightarrow S_n$, we regard $\mathcal{Z}_n$ as an S_n -analytic space. There is an exact sequence

$$\epsilon\mathcal{O}_{Z_n} \xrightarrow{d} \Omega_{Z_n/S_n}^1 \otimes_{A_n[\epsilon]} A_n \rightarrow \Omega_{Z_n/S_n}^1 \rightarrow 0.$$

Here d is defined by $d(\epsilon f) := f d\epsilon$ for $\epsilon f \in \epsilon\mathcal{O}_{Z_n}$. By using the fact that Z is reduced, one can show that this d is an injection; hence we have an element of $\text{Ext}^1(\Omega_{Z_n/S_n}^1, \epsilon\mathcal{O}_{Z_n})$. Since $\mathcal{Z}_n$ is flat over $S_n[\epsilon]$, $\epsilon\mathcal{O}_{Z_n} \cong \mathcal{O}_{Z_n}$.

Conversely, let there be an element of $\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \cong \text{Ext}^1(\Omega_{Z_n/S_n}^1, \epsilon\mathcal{O}_{Z_n})$. This element gives an $\mathcal{O}_{Z_n}$ -module extension of Ω_{Z_n/S_n}^1 by $\epsilon\mathcal{O}_{Z_n}$. We consider this extension as an A_n -module extension. Then by the homomorphism $\text{Ext}_{A_n}^1(\Omega_{Z_n/S_n}^1, \epsilon\mathcal{O}_{Z_n}) \rightarrow \text{Ext}_{A_n}^1(\mathcal{O}_{Z_n}, \epsilon\mathcal{O}_{Z_n})$ induced by $d : \epsilon\mathcal{O}_{Z_n} \rightarrow \Omega_{Z_n/S_n}^1$,

we have the commutative diagram of exact sequences

$$\begin{array}{ccccccc}
0 & \longrightarrow & \epsilon\mathcal{O}_{Z_n} & \longrightarrow & \mathcal{G} & \longrightarrow & \mathcal{O}_{Z_n} \longrightarrow 0 \\
& & \text{id} \downarrow & & D \downarrow & & d \downarrow \\
0 & \longrightarrow & \epsilon\mathcal{O}_{Z_n} & \longrightarrow & \mathcal{F} & \xrightarrow{\phi} & \Omega_{Z_n/S_n}^1 \longrightarrow 0
\end{array} \tag{1}$$

By definition $\mathcal{G} = \{(f, g) \in \mathcal{F} \oplus \mathcal{O}_{Z_n}; \phi(f) = dg\}$. We give a ring structure to $\mathcal{G}$ by $(f, g) \times (f', g') := (g'f + gf', gg')$. By the inclusion $\epsilon\mathcal{O}_{Z_n} \rightarrow \mathcal{F}$ we regard ϵ as an element of $\mathcal{F}$. Then $\mathcal{G}$ is an $A_n[\epsilon]/\epsilon^2$ algebra by defining $\epsilon \cdot (f, g) := (g\epsilon, 0)$ for $(f, g) \in \mathcal{G}$. Now put $\mathcal{G} = \mathcal{O}_{Z_n}$. Then $(Z_n, \mathcal{O}_{Z_n})$ is an infinitesimal deformation of $(Z_n, \mathcal{O}_{Z_n})$ over $S_n[\epsilon]$.

Remark. Another approach to prove (iii-a) is to use the *cotangent complex* and the associated tangent cohomology (cf. [L-S], [Pa], [Bi, §. 5]). Let L_{Z_n/S_n} be the cotangent complex for $Z_n \rightarrow S_n$. We put

$$T_{Z_n/S_n}^i := \underline{\text{Ext}}^i(L_{Z_n/S_n}, \mathcal{O}_{Z_n})$$

and

$$\mathbf{T}_{Z_n/S_n}^i := \mathbf{Ext}^i(L_{Z_n/S_n}, \mathcal{O}_{Z_n}).$$

By the definition of the cotangent complex, one has $T_{Z_n/S_n}^0 \cong T_{Z_n/S_n}$. Moreover, if Z is reduced, then the argument of [Ar 2, Prop. 6.1] shows that $T_{Z_n/S_n}^1 \cong \underline{\text{Ext}}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n})$. Since there is a natural map of complexes $L_{Z_n/S_n} \rightarrow \Omega_{Z_n/S_n}^1$, we get the maps

$$\underline{\text{Ext}}^i := \underline{\text{Ext}}^i(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \rightarrow T_{Z_n/S_n}^i$$

and

$$\text{Ext}^i := \text{Ext}^i(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \rightarrow \mathbf{T}_{Z_n/S_n}^i.$$

Therefore, we have a commutative diagram of local-to-global exact sequences:

$$\begin{array}{ccccccc}
H^1(T_{Z_n/S_n}) & \longrightarrow & \text{Ext}^1 & \longrightarrow & H^0(\underline{\text{Ext}}^1) & \longrightarrow & H^2(T_{Z_n/S_n}) \\
\downarrow & & \downarrow & & \downarrow & & \downarrow \\
H^1(T_{Z_n/S_n}^0) & \longrightarrow & \mathbf{T}_{Z_n/S_n}^1 & \longrightarrow & H^0(T_{Z_n/S_n}^1) & \longrightarrow & H^2(T_{Z_n/S_n}^0)
\end{array} \tag{2}$$

By this diagram we see that $\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \cong \mathbf{T}_{Z_n/S_n}^1$ when Z is reduced.

Next let us consider a locally trivial deformation $Z_n \rightarrow S_n$ of Z . Locally trivial deformations are controlled by $H^1(Z, T_{Z_n/S_n})$ instead of Ext^1 :

(iii-b): *There is a one-to-one correspondence between $H^1(Z, T_{Z_n/S_n})$ and the set of equivalence classes of locally trivial deformations of Z_n over $S_n[\epsilon]$.*

First let us take a Stein open cover $\{V_i\}_{i \in I}$ of Z_n . Then each locally trivial deformation $\mathcal{Z}_n$ of Z_n is constructed by patching $V_i \times_{S_n} S_n[\epsilon]$ together in such a way that, over S_n , it gives the original Z_n . Now the situation is the same as the usual deformation of a non-singular variety. By the same argument as [SGA 1], Expose.III, 6, we have the above correspondence.

(iv) Assume that Z is a $\mathbf{Q}$ -factorial projective symplectic variety with terminal singularities. By (ii), $\text{codim}(\Sigma \subset Z) \geq 4$, where $\Sigma := \text{Sing}(Z)$. We define A_n and S_n as in (iii). Suppose that $Z_n \rightarrow S_n$ is an infinitesimal deformation of Z over S_n . Let U be the regular part of Z . We put $U_n := Z_n|_U$.

Lemma 1. *The natural map*

$$H^1(Z, T_{Z_n/S_n}) \rightarrow \text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n})$$

is a bijection.

Proof. It suffices to show that the map

$$\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \rightarrow H^0(Z, \underline{\text{Ext}}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}))$$

is the zero map. Take a point $p \in \Sigma$. Let $Z(p)$ be a small Stein open neighborhood of $p \in Z$ and set $U(p) := Z(p) \cap U$. We put $Z_n(p) := Z_n|_{Z(p)}$ and $U_n(p) := U_n|_{U(p)}$. We prove that

$$\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \rightarrow H^0(Z_n(p), \underline{\text{Ext}}^1(\Omega_{Z_n(p)/S_n}^1, \mathcal{O}_{Z_n(p)}))$$

is the zero map. Since Z is Cohen-Macaulay and $\text{codim}(\Sigma \subset Z) \geq 3$, by [Ko-Mo, 12.5.6], one has

$$\text{Ext}^1(\Omega_{Z_n/S_n}^1, \mathcal{O}_{Z_n}) \cong H^1(U_n, T_{U_n/S_n})$$

and

$$H^0(Z_n(p), \underline{\text{Ext}}^1(\Omega_{Z_n(p)/S_n}^1, \mathcal{O}_{Z_n(p)})) \cong H^1(U_n(p), T_{U_n(p)/S_n}).$$

Since the symplectic 2-form ω extends to $\omega_n \in H^0(U_n, \Omega_{U_n/S_n}^2)$ by [Na], we can identify T_{U_n/S_n} with Ω_{U_n/S_n}^1 by this relative 2-form. Now we only have to prove that

$$H^1(U_n, \Omega_{U_n/S_n}^1) \rightarrow H^1(U_n(p), \Omega_{U_n(p)/S_n}^1)$$

is the zero map. Let us consider the Hodge spectral sequences

$$H^j(U_n, \Omega_{U_n/S_n}^i) \Rightarrow H^{i+j}(U, A_n)$$

and

$$H^j(U_n(p), \Omega_{U_n(p)/S_n}^i) \Rightarrow H^{i+j}(U(p), A_n),$$

where A_n is the constant sheaf with coefficient A_n . In the latter spectral sequence, we have a decreasing filtration $0 \subset F_{loc}^2 \subset F_{loc}^1 \subset F_{loc}^0 = H^2(U(p), A_n)$. By the depth argument, $E_1^{0,1} := H^1(U_n(p), \mathcal{O}_{U_n(p)}) = 0$ because Z is Cohen-Macaulay and $\text{codim}(\Sigma \subset Z) \geq 3$. So there is an injection $Gr_{F_{loc}}^1(H^2(U(p), A_n)) \rightarrow H^1(U_n(p), \Omega_{U_n(p)/S_n}^1)$. In particular, we have a map $F_{loc}^1 \rightarrow H^1(U_n(p), \Omega_{U_n/S_n}^1)$. On the other hand, the first spectral sequence degenerates at E_1 terms for $i + j = 2$ (cf. [Na 1], Lemma 2.7) because $\text{codim}(\Sigma \subset Z) \geq 4$. Hence, for the decreasing filtration $F^\cdot$ of $H^2(U, A_n)$, $Gr_F^1(H^2(U, A_n)) = H^1(U_n, \Omega_{U_n/S_n}^1)$. In particular, we have a *surjection* $F^1 \rightarrow H^1(U_n, \Omega_{U_n/S_n}^1)$. The inclusion map $U(p) \rightarrow U$ induces a map of the second cohomologies, say $\iota(p)$

$$\iota(p) : H^2(U, A_n) \rightarrow H^2(U(p), A_n).$$

This map preserves the filtrations $\{F^\cdot\}$ and $\{F_{loc}^\cdot\}$; hence it induces a map $F^1 \rightarrow F_{loc}^1$. By the commutative diagram

$$\begin{array}{ccc} F^1 & \longrightarrow & H^1(U_n, \Omega_{U_n/S_n}^1) \\ \downarrow & & \downarrow \\ F_{loc}^1 & \longrightarrow & H^1(U_n(p), \Omega_{U_n/S_n}^1) \end{array}$$

we only have to prove that the map $\iota(p)$ is the zero map. Take a resolution $\nu : \tilde{Z} \rightarrow Z$ such that $\nu^{-1}(U) \cong U$. Put $\tilde{Z}(p) := \nu^{-1}(Z(p))$. Let us consider the commutative diagram

$$\begin{array}{ccc} H^2(\tilde{Z}, A_n) & \rightarrow & H^2(U, A_n) \\ \downarrow & & \downarrow \\ H^2(\tilde{Z}(p), A_n) & \rightarrow & H^2(U(p), A_n). \end{array}$$

Here note that $H^2(\cdot, A_n) = H^2(\cdot, \mathbf{C}) \otimes_{\mathbf{C}} A_n$. Since Z is $\mathbf{Q}$ -factorial, $\text{im}[H^2(\tilde{Z}, A_n) \rightarrow H^2(\tilde{Z}(p), A_n)]$ is generated by $[E_i]$'s as an A_n module by (i), where E_i are exceptional divisors of ν . Therefore, the composite

$$H^2(\tilde{Z}, A_n) \rightarrow H^2(\tilde{Z}(p), A_n) \rightarrow H^2(U(p), A_n)$$

is the zero map. On the other hand, the map $H^2(\tilde{Z}, A_n) \rightarrow H^2(U, A_n)$ is a surjection by an argument in [Na, Proposition 9](cf. (b), p.143) because $\text{codim}(\Sigma \subset Z) \geq 4$. Now $\iota(p)$ is the zero map by the commutative diagram. Q.E.D.

Let us start the proof of Main Theorem. Let A be an Artinian local $\mathbf{C}$ algebra and put $S := \text{Spec}(A)$. Recall that a deformation $\mathcal{Z} \rightarrow S$ of Z is locally trivial if, for any Stein open set $V \subset Z$, $\mathcal{Z}|_V \cong V \times S$. We can introduce a *locally trivial* deformation functor of Z

$$D_{lt} : (\text{Art})_{\mathbf{C}} \rightarrow (\text{Set})$$

as a sub-functor of the usual deformation functor D of Z . One can check that D_{lt} has a pro-representable hull $(R_{lt}, \mathcal{Z}_{R_{lt}})$ in the sense of Schlessinger [Sch]. Here R_{lt} is a complete local $\mathbf{C}$ algebra and $\mathcal{Z}_{R_{lt}}$ is a certain projective system of infinitesimal deformations of Z . Moreover, by the definition of D_{lt} , one can construct R_{lt} as a suitable quotient of the hull R of the usual deformation functor D . D has T^1 -lifting property as noted in [Na] (for the relationship between T^1 -lifting property and unobstructedness of deformations, see [Ka], [Na 3]). Hence, by Lemma 1 and (iii-b), we see that D_{lt} also has T^1 -lifting property. This implies that R and R_{lt} are both complete regular local ring over $\mathbf{C}$. Since their cotangent spaces coincide again by Lemma 1, we conclude that $R \cong R_{lt}$. This implies that any flat deformation of Z is locally trivial.

3 Applications

In this section we shall generalize Theorem 2.2 of [Na 1] to the case of $\mathbf{Q}$ -factorial terminalizations (cf. Theorem 1) and apply Main Theorem and Theorem 1 to get Corollaries 1 and 2. Propositions 1 and 2 are preliminaries for the proof of Theorem 1.

Let us recall some generalities on deformation theory. Let Y be a compact complex space. Then it has been proved by [Gr], [Do], [Pa] and others that, there is a Kuranishi space $0 \in \text{Def}(Y)$ and a semi-universal deformation $f : \mathcal{Y} \rightarrow \text{Def}(Y)$ of Y with $f^{-1}(0) = Y$. Assume that $h : Y \rightarrow Z$ is a (proper) map of compact complex spaces such that $R^1 h_* \mathcal{O}_Y = 0$ and $h_* \mathcal{O}_Y = \mathcal{O}_Z$. Let $\mathcal{Y} \rightarrow \text{Def}(Y)$ and $\mathcal{Z} \rightarrow \text{Def}(Z)$ be the semi-universal deformations. Then one can construct two maps $h_* : \text{Def}(Y) \rightarrow \text{Def}(Z)$ and $H_* : \mathcal{Y} \rightarrow \mathcal{Z}$ in such

a way that the following diagram commutes (cf. [Ko-Mo, 11.4])

$$\begin{array}{ccc}
 \mathcal{Y} & \xrightarrow{H_*} & \mathcal{Z} \\
 \downarrow & & \downarrow \\
 \mathrm{Def}(Y) & \xrightarrow{h_*} & \mathrm{Def}(Z)
 \end{array} \tag{3}$$

For the convenience of readers, we shall give a sketch of the construction of H_* and h_* according to [Ko-Mo, 11.4]. First note that H_* and h_* exist in the formal category by [Wa]. We want to construct them in the analytic category. Let $G \subset Y \times Z$ be the graph of h . We shall find a graph $\mathcal{G} \subset \mathcal{Y} \times \mathcal{Z}$ in such a way that $\mathcal{G} \cap (Y \times Z) = G$. Let us consider the component D of the relative Douady space of $\mathcal{Y} \times \mathcal{Z} / \mathrm{Def}(Y) \times \mathrm{Def}(Z)$ containing $[G]$. The projection morphism $D \rightarrow \mathrm{Def}(Y)$ has a formal section defined by the formal contraction morphism around $0 \in \mathrm{Def}(Y)$. Then, by [Ar 1, 1.5], one can find an analytic section. Note that, when h is a birational map, $(H_*)_t : Y_t \rightarrow Z_{h_*(t)}$ is a birational map for $t \in \mathrm{Def}(Y)$.

Proposition 1. *Let Z be a projective symplectic variety. Assume that $h : Y \rightarrow Z$ is a $\mathbf{Q}$ -factorial terminalization, that is, h is a birational morphism and Y is a $\mathbf{Q}$ -factorial projective symplectic variety with terminal singularities. Then there is a deformation of $h : \mathcal{Y}_\Delta \xrightarrow{h} \mathcal{Z}_\Delta \rightarrow \Delta$ over a disc such that for $t \in \Delta \setminus 0$, $\tilde{h}_t : Y_t \rightarrow Z_t$ is an isomorphism.*

Proof. Denote by $(T, 0)$ the Kuranishi space of Y and let $p : \mathcal{Y} \rightarrow T$ be the semi-universal family over T . By [Na 1], Proposition (2.5), T is smooth. Take a projective resolution $\pi : \tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$ in such a way that π is an isomorphism outside $\mathrm{Sing}(\mathcal{Y})$. For a general $t \in T$, $\pi_t : \tilde{Y}_t \rightarrow Y_t$ is a resolution. Moreover, $\pi_t^{-1}(U_t) \cong U_t$ for the regular locus U_t of Y_t , since $U_t \cap \mathrm{Sing}(\mathcal{Y}) = \emptyset$. Note that $\tilde{Y}_t$ is not, in general, a projective variety, but a Kähler manifold. We shall prove that one can deform Y_t further to a variety $Y_{t'}$ so that it contains no curves. Since there is a map of Kuranishi spaces $h_* : T \rightarrow \mathrm{Def}(Z)$ as noted above, one has a birational morphism $Y_{t'} \rightarrow Z_{h_*(t')}$. But, $Y_{t'}$ does not have any curve; hence this birational morphism should be an isomorphism (cf. [Na 1], claim 3, p. 618). Since Y_t is a small deformation of a symplectic variety Y , Y_t is again symplectic by [Na]. A symplectic singularity is a canonical singularity (cf. [Be]). A canonical singularity is a rational singularity by [El]. Therefore, Y_t has only rational singularities. Since Y_t has only rational singularities, the natural map $H^2(Y_t, \mathbf{C}) \rightarrow H^2(\tilde{Y}_t, \mathbf{C})$ is an

injection of mixed Hodge structures ([De]). Hence, by the strict compatibility of weight filtrations, $H^2(Y_t, \mathbf{C})$ admits a pure Hodge structure of weight 2. Let

$$H^2(Y_t, \mathbf{C}) = H^{2,0}(Y_t) \oplus H^{1,1}(Y_t) \oplus H^{0,2}(Y_t)$$

be the Hodge decomposition. Recall again that there is an injection of pure Hodge structures $H^2(Y_t, \mathbf{C}) \rightarrow H^2(\tilde{Y}_t, \mathbf{C})$. Put $n := \dim Y$ and consider the perfect pairing

$$\langle , \rangle : H^{2n-2}(\tilde{Y}_t, \mathbf{C}) \times H^2(\tilde{Y}_t, \mathbf{C}) \rightarrow H^{2n}(\tilde{Y}_t, \mathbf{C}) = \mathbf{C}.$$

Set

$$V_t := H^2(Y_t, \mathbf{C})^\perp \subset H^{2n-2}(\tilde{Y}_t, \mathbf{C}).$$

Here we regard $H^2(Y_t, \mathbf{C})$ as a subspace of $H^2(\tilde{Y}_t, \mathbf{C})$.

Claim 1. *Every element of V_t is of type $(n-1, n-1)$.*

Proof. If we put

$$V_t^{n-1, n-1} := V_t \cap H^{n-1}(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-1}),$$

then we see that $V_t^{n-1, n-1} = H^{1,1}(Y_t)^\perp \subset H^{n-1}(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-1})$ for the Serre pairing

$$\langle , \rangle : H^{n-1}(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-1}) \times H^1(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^1) \rightarrow H^n(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^n) = \mathbf{C}.$$

Since $h^{n-1, n-1}(\tilde{Y}_t) = h^{1,1}(\tilde{Y}_t)$, $\dim V_t^{n-1, n-1} = h^{1,1}(\tilde{Y}_t) - h^{1,1}(Y_t)$. On the other hand, $\dim V_t = b_2(\tilde{Y}_t) - b_2(Y_t)$. We shall prove that $\dim V_t = \dim V_t^{n-1, n-1}$. In order to do this, it suffices to show that all elements of $H^2(\tilde{Y}_t, \mathbf{C})$ of type $(2,0)$ and of type $(0,2)$ are mapped to zero by the map $\phi : H^2(\tilde{Y}_t, \mathbf{C}) \rightarrow H^0(Y_t, R^2\pi_{t*}\mathbf{C})$. In fact, since the map $H^2(Y_t) \rightarrow H^2(\tilde{Y}_t)$ is an injection of pure Hodge structures of weight 2, if these are proved, then the injection is actually an isomorphism at $(2,0)$ and $(0,2)$ parts. By the conjugation, it suffices to show that $\phi(\alpha) = 0$ for an element $\alpha \in H^2(\tilde{Y}_t)$ of type $(2,0)$. We shall prove that, for every point $x \in Y_t$, $\phi(\alpha)_x = 0$ in $(R^2(\pi_t)_*\mathbf{C})_x$. Let $\nu : W \rightarrow \tilde{Y}_t$ be a projective birational morphism such that W is smooth and $D := (\pi_t \circ \nu)^{-1}(x)$ is a simple normal crossing divisor of W . Put $h := \pi_t \circ \nu$. Since $R^1\nu_*\mathbf{C} = 0$, $R^2(\pi_t)_*\mathbf{C}$ injects to $R^2h_*\mathbf{C}$; hence we have to check that α is sent to zero by the composite

$$H^2(\tilde{Y}_t, \mathbf{C}) \rightarrow H^2(W, \mathbf{C}) \rightarrow (R^2h_*\mathbf{C})_x (= H^2(D, \mathbf{C})).$$

We call this composite ψ . Then ψ preserves Hodge filtrations of $H^2(\tilde{Y}_t)$ and $H^2(D)$. Hence it induces $Gr_F^2(H^2(\tilde{Y}_t)) \rightarrow Gr_F^2(H^2(D)) \subset H^2(D)$. Since $\alpha \in Gr_F^2(H^2(\tilde{Y}_t))$, $\psi(\alpha) \in Gr_F^2(H^2(D))$. But, since Y_t has rational singularities at x , $Gr_F^2(H^2(D)) = H^0(D, \hat{\Omega}_D^2) = 0$ by [Na 1, Lemma (1.2)], where $\hat{\Omega}_D^2 := \Omega_D^2 / (\text{torsion})$. Therefore, $\dim V_t = \dim V_t^{n-1, n-1}$. Q.E.D.

The families

$$\tilde{\mathcal{Y}} \rightarrow \mathcal{Y} \rightarrow T$$

induces the Kodaira-Spencer maps

$$\rho_{\tilde{Y}_t} : T_t(T) \rightarrow H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t}),$$

$$\rho_{Y_t} : T_t(T) \rightarrow \text{Ext}^1(\Omega_{Y_t}^1, \mathcal{O}_{Y_t}).$$

Since $\mathcal{Y} \rightarrow T$ is versal at $t \in T$, ρ_{Y_t} is surjective. Define a map

$$d : H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t}) \rightarrow \text{Ext}^1(\Omega_{Y_t}^1, \mathcal{O}_{Y_t})$$

as the *differential* of the map $\text{Def}(\tilde{Y}_t) \rightarrow T$. We then have $d \circ \rho_{\tilde{Y}_t} = \rho_{Y_t}$. Let $U_t \subset Y_t$ be the regular locus of Y_t . Then $\text{Ext}^1(\Omega_{Y_t}^1, \mathcal{O}_{Y_t}) \cong H^1(U_t, \Theta_{U_t})$. By this identification, the map d coincides with the restriction map $H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t}) \rightarrow H^1(U_t, \Theta_{U_t})$. A symplectic 2-form on Y_t extends to a holomorphic 2-form on $\tilde{Y}_t$. By this 2-form, we have a map $H^1(\Theta_{\tilde{Y}_t}) \rightarrow H^1(\Omega_{\tilde{Y}_t}^1)$. We put

$$T_1 := \text{im}[T_t(T) \rightarrow H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t})],$$

$$T_2 := \text{im}[T_t(T) \rightarrow H^1(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^1)].$$

Claim 2.

$$\dim T_2 \geq \dim H^1(U_t, \Omega_{U_t}^1).$$

Proof. Let us consider the composite

$$T_t(T) \rightarrow H^1(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^1) \rightarrow H^1(U_t, \Omega_{U_t}^1).$$

By the symplectic 2-form, we identify $H^1(U_t, \Theta_{U_t})$ with $H^1(U_t, \Omega_{U_t}^1)$. Then the composite above coincides with the Kodaira-Spencer map ρ_{Y_t} , which is surjective.

Look at the pairing

$$(\ , \) : H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t}) \times H^{n-1}(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-1}) \rightarrow H^n(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-2})$$

Claim 3. *In the pairing above, $(T_1, V_t) = 0$.*

Proof. For $\zeta \in H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t})$, take the corresponding infinitesimal deformation $\tilde{Y}_{t,\epsilon} \rightarrow \text{Spec } \mathbf{C}[\epsilon]$. Choose an element $l \in H^{2n-2}(\tilde{Y}_t, \mathbf{C})$ of type $(n-1, n-1)$. Then (ζ, l) is an obstruction for the class l to remain of type $(n-1, n-1)$ under the infinitesimal deformation. Now the claim follows from Claim 1 and the fact that $R^2p_*\mathbf{C}_Y$ is a local system on T (cf. [Na]¹).

Since T_2 is the image of T_1 by the map $H^1(\Theta_{\tilde{Y}_t}) \rightarrow H^1(\Omega_{\tilde{Y}_t}^1)$, we have $\langle T_2, V_t \rangle = 0$ by the commutative diagram of the pairing maps:

$$\begin{array}{ccc} (,) : H^1(\tilde{Y}_t, \Theta_{\tilde{Y}_t}) \times H^{n-1}(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-1}) & \rightarrow & H^n(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-2}) \\ \downarrow & & \downarrow \\ \langle , \rangle : H^1(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^1) \times H^{n-1}(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^{n-1}) & \rightarrow & H^n(\tilde{Y}_t, \Omega_{\tilde{Y}_t}^n). \end{array}$$

Note that $\text{codim}(V_t \subset H^{n-1, n-1}(\tilde{Y}_t)) = h^{1,1}(Y_t)$. On the other hand, since $h^{1,1}(Y_t) = h^1(U_t, \Omega_{U_t}^1)$ (cf. [Na]²), we see, by Claim 2, that $T_2 = (V_t)^\perp$ with respect to the pairing $\langle , \rangle$ above.

Now let C be a connected curve on $\tilde{Y}_t$ such that $\pi_t(C)$ is not a point. Since $[C] \notin V_t$, we see that $[C]^\perp \cap T_2$ is a codimension 1 subspace of T_2 by the perfectness of $\langle , \rangle$. In particular, we can find an element $\zeta \in T_1$ such that $(\zeta, [C]) \neq 0$. The $(n-1, n-1)$ classes of curves on $\tilde{Y}_t$ constitute a countable subset of $H^{n-1, n-1}(\tilde{Y}_t)$. Hence, if we take a generic deformation $\tilde{Y}_{t'}$ of $\tilde{Y}_t$, then any effective 1-cycle on $\tilde{Y}_t$ which is not contained in a fiber of π_t cannot deform sideways. Since $\tilde{\mathcal{Y}} \rightarrow T$ is a Kaehler morphism, every

¹In [Na], this has been proved under certain additional conditions: $h^0(Y_{\text{reg}}, \Omega_{Y_{\text{reg}}}^2) = 1$, $h^1(Y, \mathcal{O}_Y) = 0$. However, these conditions are not essential. Here we give a more direct proof to this fact by using our Main Theorem. Since $\mathcal{Y} \rightarrow T$ is proper, if we choose T small enough, then there is an open covering $\{\mathcal{V}_i\}_{i \in I}$ of $\mathcal{Y}$ with (analytic) trivializations over T : $\mathcal{V}_i \cong V_i \times T$. Here $V_i := \mathcal{Y} \cap \mathcal{V}_i$. Let us take a canonical stratification $Y = \coprod_j Y_j$ of Y into locally closed smooth subsets such that $\bar{Y}_j = \text{Sing}(\bar{Y}_{j-1})$ and $Y_0 = Y_{\text{reg}}$. Fix a nowhere vanishing C^∞ -vector field ζ of T . One can lift ζ to a C^∞ -vector field ζ_i of $V_i \times T$ in such a way that ζ_i induces a vector field of each $(Y_j \cap V_i) \times T$ for each stratum Y_j . By using the partition of unity, one can glue together $\{\zeta_i\}$ and get a globally defined vector field $\tilde{\zeta}$ of $\mathcal{Y}$. This vector field gives us a C^∞ -trivialization of $\mathcal{Y}$ over T . Hence $R^ip_*\mathbf{C}$ are constant sheaves.

²By the argument of the footnote (1) above, we see that $h^2(Y) = h^2(Y_t)$ and $h^2(U) = h^2(U_t)$ with $U := Y_{\text{reg}}$. Since Y is $\mathbf{Q}$ -factorial, one has $H^2(Y) \cong H^2(U)$; hence $H^2(Y_t) \cong H^2(U_t)$. Now we get the result by the footnote (1) in p. 21 of the eprint version of [Na] (math.AG/0010114).

irreducible component of the Douady space $D(\tilde{\mathcal{Y}}/T)$ parametrizing curves is proper over T ([Fu 2]). Therefore, all components dominating T parametrize the curves which are contracted to points by $\tilde{\mathcal{Y}} \rightarrow \mathcal{Y}$. By the countability of the Douady space ([Fu 3]), we now see that, for a generic deformation $\tilde{Y}_{t'}$ of $\tilde{Y}_t$, any holomorphic curve C on $\tilde{Y}_{t'}$ is $\pi_{t'}$ -exceptional for $\pi_{t'} : \tilde{Y}_{t'} \rightarrow Y_{t'}$, that is, $\pi_{t'}(C)$ is a point. Then, the variety $Y_{t'}$ has no holomorphic curves. In fact, if there is a curve C , then one can find a curve D on $\tilde{Y}_{t'}$ such that $\pi_{t'}(D) = C$ by using the Chow lemma [Hi], which is a contradiction.

Let $h : Y \rightarrow Z$ be a $\mathbf{Q}$ -factorial terminalization of a projective symplectic variety Z . We put $\Sigma := \text{Sing}(Z)$. There is a closed subset Σ_0 of Σ with $\text{codim}(\Sigma_0 \subset Z) \geq 4$ such that Z is locally a trivial family of a rational double point along $\Sigma \setminus \Sigma_0$. In the remainder, we put $U := Z \setminus \Sigma_0$ and $V := Y \setminus \text{Sing}(Y)$. Note that $h^{-1}(U) \subset V$. We denote by n the dimension of Z . In this situation, we have the following generalization of [Na 1], Proposition (2.1).

Proposition 2³. *There is a commutative diagram*

$$\begin{array}{ccc} \text{Ext}^1(\Omega_Y^1, \mathcal{O}_Y) & \rightarrow & H^1(h^{-1}(U), \Theta_{h^{-1}(U)}) \\ \downarrow & & \downarrow \\ \text{Ext}^1(\Omega_Z^1, \mathcal{O}_Z) & \rightarrow & \text{Ext}^1(\Omega_U^1, \mathcal{O}_U), \end{array}$$

where horizontal maps are both isomorphisms.

Proof. In the diagram, each space corresponds to the set of first order deformations of Y , $h^{-1}(U)$, Z or U , respectively. A first order deformation of Y (resp. $h^{-1}(U)$) induces that of Z (resp. U) (cf. [Na 1], p.614). Each vertical map is nothing but this correspondence. On the other hand, the horizontal maps are natural ones induced by the restriction. The second horizontal map (at the bottom) is an isomorphism by [Ko-Mo, 12.5.6] because $\text{codim}(\Sigma_0 \subset Z) \geq 3$ and Z is Cohen-Macaulay. By the same reason, $\text{Ext}^1(\Omega_Y^1, \mathcal{O}_Y) \cong H^1(V, \Theta_V)$. We only have to prove that the first horizontal map is an isomorphism. Set $F := Y \setminus h^{-1}(U)$ and $F^0 := V \setminus h^{-1}(U)$. Note that $\dim F \leq n - 2$ since h is a crepant partial resolution of a symplectic variety ([Na 1], Corollary (1.15); see also [Na 2], footnote, p.1). Let us consider the exact sequence of cohomology with coefficient $\mathbf{C}$

$$H_{F^0}^2(V) \rightarrow H^2(V) \rightarrow H^2(h^{-1}(U)) \rightarrow H_{F^0}^3(V).$$

³ $\mathbf{Q}$ -factoriality of Y is not necessary for this proposition

Let $j : h^{-1}(U) \rightarrow V$ and $i : F^0 \rightarrow V$ be the inclusion maps. Then the dual of this sequence coincides with the exact sequence of cohomology with compact support:

$$H_c^{2n-3}(F^0) \rightarrow H_c^{2n-2}(h^{-1}(U)) \rightarrow H_c^{2n-2}(V) \rightarrow H_c^{2n-2}(F^0)$$

induced by the exact sequence (cf. [Ha, II, Ex.1.19(c)])

$$0 \rightarrow j_!(\mathbf{C}_{h^{-1}(U)}) \rightarrow \mathbf{C}_V \rightarrow i_*(\mathbf{C}_{F^0}) \rightarrow 0.$$

Here we have $H_c^{2n-3}(F^0) = H_c^{2n-2}(F^0) = 0$. In fact, the first terms and the third terms of the next exact sequences vanish (cf. [L]) since $\dim_{\mathbf{C}} F \leq n-2$ and $\dim_{\mathbf{C}}(F \setminus F^0) \leq n-3$:

$$H^{2n-3}(F \setminus F^0) \rightarrow H_c^{2n-2}(F^0) \rightarrow H^{2n-2}(F)$$

and

$$H^{2n-4}(F \setminus F^0) \rightarrow H_c^{2n-3}(F^0) \rightarrow H^{2n-3}(F).$$

As a consequence, the map

$$H^2(V) \rightarrow H^2(h^{-1}(U))$$

is an isomorphism. Moreover, this is a morphism of mixed Hodge structures. Since Y is a symplectic variety with terminal singularities, $\text{codim}(\text{Sing}(Y) \subset Y) \geq 4$; hence $H^2(V)$ is equipped with a pure Hodge structure of weight 2 with $\text{Gr}_F^1(H^2(V)) = H^1(\Omega_V^1)$ (cf. e-print version math.AG/0010114 of [Na], footnote at p.21). Let us examine the mixed Hodge structure on $H^2(h^{-1}(U))$. Let $\nu : \tilde{Y} \rightarrow Y$ be a projective birational morphism from a non-singular variety $\tilde{Y}$ to Y such that $\nu^{-1}(h^{-1}(U)) \cong h^{-1}(U)$ and such that $\text{Exc}(h \circ \nu)$ is a normal crossing variety of $\tilde{Y}$. We put $E' := \tilde{Y} \setminus (h \circ \nu)^{-1}(U)$. By [De], $\text{Gr}_F^1(H^2(h^{-1}(U))) = H^1(\tilde{Y}, \Omega_{\tilde{Y}}^1(\log E'))$. Let us consider the exact sequence of local cohomology

$$H_{E'}^1(\Omega_{\tilde{Y}}^1(\log E')) \rightarrow H^1(\Omega_{\tilde{Y}}^1(\log E')) \rightarrow H^1(h^{-1}(U), \Omega_{h^{-1}(U)}^1) \rightarrow H_{E'}^2(\Omega_{\tilde{Y}}^1(\log E')).$$

The first term and the fourth term both vanish. This follows from the same argument as the proof of [Na 1], Proposition (2.1): by [Na 1, claim 2, p.616], $R^k(h \circ \nu)_* \Omega_{\tilde{Y}}^{n-1}(\log E')(-E') = 0$ for $k \geq 2$. Then by taking the duals of the first term and the fourth term, we get the conclusion since $\text{codim}(\Sigma_0 \subset Z) \geq 4$ (cf. [Na 1, p.615]). As a consequence, we see that

$$\text{Gr}_F^1(H^2(h^{-1}(U))) = H^1(h^{-1}(U), \Omega_{h^{-1}(U)}^1).$$

Since the natural map

$$H^1(V, \Omega_V^1) \rightarrow H^1(h^{-1}(U), \Omega_{h^{-1}(U)}^1)$$

coincides with the map

$$\mathrm{Gr}_F^1(H^2(V)) \rightarrow \mathrm{Gr}_F^1(H^2(h^{-1}(U))),$$

we conclude that it is an isomorphism. Finally, by the symplectic 2-form on V , this map is identified with

$$H^1(V, \Theta_V) \rightarrow H^1(h^{-1}(U), \Theta_{h^{-1}(U)}).$$

Theorem 1. *Let $h : Y \rightarrow Z$ be a $\mathbf{Q}$ -factorial terminalization of a projective symplectic variety Z . Then the Kuranishi spaces $\mathrm{Def}(Y)$ and $\mathrm{Def}(Z)$ are both smooth of the same dimension. The natural map (cf. [Na 1]) $h_* : \mathrm{Def}(Y) \rightarrow \mathrm{Def}(Z)$ is a finite covering. Moreover, Z has a flat deformation to a (non-projective) symplectic variety Z_t which is, at the same time, a small deformation of Y .*

Proof. When Y is non-singular, it is nothing but Theorem (2.2) of [Na 1]. In a general case, we modify the proof of [Na 1] in the following way. First, Corollary (1.5) of [Na 1] holds true when $\tilde{X}$ is a $\mathbf{Q}$ -factorial terminalization of X (here the notation being the same as [Na 1]). Next, Proposition (1.6) of [Na 1] follows from a more general fact recently proved in [K]. Finally, we have to replace Proposition (2.1) of [Na 1] by Proposition 2 and replace Claim 3 of [Na 1](p.618) by Proposition 1.

Corollary 1. *Let $h : Y \rightarrow Z$ and $h' : Y' \rightarrow Z$ be two $\mathbf{Q}$ -factorial terminalizations of a projective symplectic variety Z . Assume that Y is non-singular. Then Y' is also non-singular.*

Proof. Since h is a symplectic resolution, the Kuranishi space $\mathrm{Def}(Z)$ of Z is non-singular by [Na 1, Theorem (2.2)]. Moreover, a general point of $\mathrm{Def}(Z)$ parametrizes a non-singular variety. Assume now that Y' is singular. By Proposition 1, there is a point $t \in \mathrm{Def}(Z)$ parametrizing a suitable deformation Y'_t of Y' . By Main Theorem, Y' does not have any smoothing via deformation; hence Y'_t does not have, too. This contradicts the fact that a general point of $\mathrm{Def}(Z)$ parametrizes a non-singular variety.

Corollary 2. *Let Z be a projective symplectic variety. Assume that Z has a $\mathbf{Q}$ -factorial terminalization $h : Y \rightarrow Z$. Then the following are equivalent.*

- (1) Z is smoothable by a flat deformation;
- (2) Y is non-singular.

Proof. (1) $\Rightarrow$ (2): Suppose that Y is singular. Take a small deformation Y_t of Y . Then Y_t is not smoothable by any flat deformation by Main Theorem. Applying Proposition 1 to $h : Y \rightarrow Z$, we see that $Z_{h_*(t)}(\cong Y_t)$ is also non-smoothable. But, by Theorem 1, $\text{Def}(Z)$ is non-singular and hence irreducible. This contradicts the assumption (1).

(2) $\Rightarrow$ (1): This is nothing but Proposition 1.

References

- [Ar 1] Artin, M.: On the solutions of analytic equations, *Invent. Math.* **5** (1968), 277-291.
- [Ar 2] Artin, M.: Lectures on deformations of singularities, Tata Inst. Fund. Res. Bombay (1976)
- [Be] Beauville, A.: Symplectic singularities, *Invent. Math.* **139** (2000), no. 3, 541-549.
- [Bi] Bingener, J.: Offenheit der Versalität in der analytischen Geometrie, *Math. Z.* **173** (1980), 241-281
- [De] Deligne, P.: Théorie de Hodge II, *Publ.Math. IHES.* **40** (1971), 5-58; III, *ibid.* **44** (1974), 5-77
- [Do] Douady, A.: Le problème des modules locaux pour les espaces C -analytiques compacts, *Ann. Sci. École Norm. Sup. (4)* **7** (1974), 569-602
- [El] Elkik, R.: Rationalité des singularités canoniques, *Invent. Math.* **64** (1981), no. 1, 1-6.
- [Fu 1] Fujiki, A.: On primitively symplectic compact Kaehler V-manifolds, in: Classification theory of algebraic and analytic manifolds, *Prog. Math.* **39** (1983), 71-250
- [Fu 2] Fujiki, A.: Closedness of the Douady spaces of compact Kähler spaces, *Publ. Res. Inst. Math. Sci.* **14** (1978/79), no. 1, 1-52.

- [Fu 3] Fujiki, A.: Countability of the Douady space of a complex space. Japan. J. Math. (N.S.) **5** (1979), no. 2, 431-447.
- [G-K] Ginzburg, V., Kaledin, D.: Poisson deformations of symplectic quotient singularities, Adv. Math. **186** (2004), 1-57
- [Gr] Grauert, H.: Der Satz von Kuranishi für kompakte komplexe Räume, Invent. Math. **25** (1974), 107-142
- [Ha] Hartshorne, R.: Algebraic Geometry, Graduate texts **49**, Springer, Heidelberg (1977)
- [Hi] Hironaka, H.: Flattening theorem in complex-analytic geometry, Amer. J. Math. **97** (1975), 503-547
- [K] Kaledin, D.: Symplectic singularities from the Poisson view point, math.AG/0310186, to appear in J. Reine Angew. Math.
- [Ka] Kawamata, Y.: Unobstructed deformations, a remark on a paper by Z. Ran, J. Algebraic Geom. **1** (1992), 183-190, Erratum, ibid, **6** (1997), 803-804
- [K-L-M] Kaledin, D., Lehn, M., Sorger, C.: Singular symplectic moduli spaces, math.AG/0504202
- [Ko-Mo] Kollár, J., Mori, S.: Classification of the three dimensional flips, J. Amer. Math. Soc. **5**(3) (1992), 533-703
- [L] Łojasiewicz, S.: Triangulation of semi-analytic sets, Ann. Sc. Norm. Super. Pisa III Ser. **18** (1964), 449-474
- [L-S] Lichtenbaum, S., Schlessinger, M.: The cotangent complex of a morphism, Trans. Amer. Math. Soc. **128** (1967), 41-70
- [Na] Namikawa, Y.: Extension of 2-forms and symplectic varieties, J. Reine Angew. Math. **539** (2001), 123-147
- [Na 1] Namikawa, Y.: Deformation theory of singular symplectic n-folds, Math. Ann. **319** (2001) 597-623
- [Na 2] Namikawa, Y.: A note on symplectic singularities, math.AG/0101028

- [Na 3] Namikawa, Y.: Calabi-Yau threefolds and deformation theory, Sugaku Exposition, **15**, vol.1, (2002), 1-29
- [O] O'Grady, K.: Desingularized moduli spaces of sheaves on a K3, J. Reine Angew. Math. **512** (1999), 49-117
- [Pa] Palamodov, V.P.: Deformations of complex spaces, Russian Math. Survey, **31-3** (1976), 129-197
- [Re] Reid, M.: Canonical 3-folds, in: Journées de Géométrie Algébrique d'Angers (1979) Sijthoff and Noordhoff, Alphen aan den Rijn, 273-310
- [Sch] Schlessinger, M.: Functors of Artin rings, Trans. Amer. Math. Soc. **130** (1968), 208-222
- [SGA 1] Grothendieck et al.: SGA 1, Lecture Note in Math. **224**, Springer (1970)
- [Sh] Shokurov, V.V.: Prelimiting flips, Proc. Steklov Inst. Math. (2003), no. 1(240), 75-213
- [Wa] Wahl, J.: Equisingular deformations of normal surface singularities I, Ann. of Math. (2) **104** (1976), no. 2, 325-356.

Yoshinori Namikawa
Departement of Mathematics, Graduate School of Science, Osaka University,
JAPAN
namikawa@math.sci.osaka-u.ac.jp